

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Parathyroid Hormone Antibody / C-Terminal, Clone: [PTH/1173], Mouse, Monoclonal NSJ-V2809IHC-7ML

Artikelname	Parathyroid Hormone Antibody / C-Terminal, Clone: [PTH/1173], Mouse, Monoclonal
Artikelnummer	NSJ-V2809IHC-7ML
Hersteller Artikelnummer	V2809IHC-7ML
Alternativnummer	NSJ-V2809IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 32-115 of human PTH was used as the immunogen for the Parathyroid Hormone antibody.
Produktbeschreibung	Epitope of this mAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood Ca ²⁺ levels by dissolving the salts in bone ...
Klonalität	Monoclonal
Klon-Bezeichnung	[PTH/1173]
UniProt	P01270

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Parathyroid Hormone antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.