

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### PTH Antibody / C-Terminal, Clone: [PTH/1175], Mouse, Monoclonal NSJ-V2811-20UG

Artikelname	PTH Antibody / C-Terminal, Clone: [PTH/1175], Mouse, Monoclonal
Artikelnummer	NSJ-V2811-20UG
Hersteller Artikelnummer	V2811-20UG
Alternativnummer	NSJ-V2811-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 32-115 of human PTH was used as the immunogen for the Parathyroid Hormone antibody.
Produktbeschreibung	Epitope of this mAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood Ca <sup>2+</sup> levels by dissolving the salts in bone ...
Klonalität	Monoclonal
Klon-Bezeichnung	[PTH/1175]
UniProt	<a href="#">P01270</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the PTH antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.