

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### PTH Antibody Cocktail / C + N-Terminal, Clone: [3H9 + PTH/1175], Mouse, Monoclonal NSJ-V2812-20UG

|                          |   |
|--------------------------|---|
| Artikelname              | PTH Antibody Cocktail / C + N-Terminal, Clone: [3H9 + PTH/1175], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V2812-20UG  |
| Hersteller Artikelnummer | V2812-20UG  |
| Alternativnummer         | NSJ-V2812-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Synthetic peptides corresponding to amino acids 1 to 34 (clone 3H9) and 32-115 (clone PTH/1175) were used as the immunogen for the PTH antibody cocktail.   |
| Produktbeschreibung      | PTH is a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood Ca <sup>2+</sup> levels by dissolving the salts in bone and preventing their renal excretion. It is ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [3H9 + PTH/1175]  |
| UniProt                  | <a href="#">P01270</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | Protein G affinity chromatography   |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT  |
| Anwendungsbeschreibung | Optimal dilution of the PTH antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |