

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Parathyroid Hormone Antibody (N-Terminal) / Rabbit Monoclonal, Clone: [PRTM1-1R] NSJ-V3765-20UG

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Parathyroid Hormone Antibody (N-Terminal) / Rabbit Monoclonal, Clone: [PRTM1-1R]  |
| Artikelnummer            | NSJ-V3765-20UG  |
| Hersteller Artikelnummer | V3765-20UG  |
| Alternativnummer         | NSJ-V3765-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A synthetic peptide from the N-terminal region of human Parathyroid hormone was used as the immunogen for the recombinant Parathyroid Hormone antibody.   |
| Produktbeschreibung      | Parathyroid hormone (PTH), parathormone or parathyrin, is secreted by the chief cells of the parathyroid glands as a polypeptide containing 84 amino acids, yet effective hormone-receptor interaction requires solely the 34-N-terminal amino acids. Whil... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PRTM1-1R]  |
| UniProt                  | <a href="#">P01270</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | Protein A affinity chromatography   |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1)   |
| Anwendungsbeschreibung | The stated application concentrations are suggested starting points. Titration of the recombinant Parathyroid Hormone antibody may be required due to differences in protocols and secondary/substrate sensitivity.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |