

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant c-Myc Antibody, Clone: [MYC2895R], Rabbit, Monoclonal NSJ-V7971-20UG

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant c-Myc Antibody, Clone: [MYC2895R], Rabbit, Monoclonal   |
| Artikelnummer            | NSJ-V7971-20UG  |
| Hersteller Artikelnummer | V7971-20UG  |
| Alternativnummer         | NSJ-V7971-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, FACS, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Amino acids 408-439 were used as the immunogen for the recombinant c-Myc antibody. The specific epitope has been localized to amino acids 410-419 (EQKLISEEDL).   |
| Produktbeschreibung      | It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This MAb shows no cross-reaction with v-myc. c-myc... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [MYC2895R]  |
| UniProt                  | <a href="#">P01106</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | Protein A affinity chromatography   |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                                      |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | ELISA (order BSA-free format for coating), Flow cytometry: 1-2ug/million cells in 0.1ml, Western blot: 2-4ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the recombinant c-Myc antibody should be determined by the researcher.                      |