

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PD1 Antibody / PDCD1, Clone: [RMP1-14], Rat, Monoclonal NSJ-V8347-20UG

| | |
|--------------------------|---|
| Artikelname | PD1 Antibody / PDCD1, Clone: [RMP1-14], Rat, Monoclonal |
| Artikelnummer | NSJ-V8347-20UG |
| Hersteller Artikelnummer | V8347-20UG |
| Alternativnummer | NSJ-V8347-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rat |
| Kategorie | Antikörper |
| Applikation | ELISA, FACS, IF, NeA |
| Spezies Reaktivität | Mouse |
| Immunogen | PDCD1 transfected BHK cells were used as the immunogen for the PD1 antibody. |
| Produktbeschreibung | PD-1 is expressed on a subset of CD4-CD8-thymocytes, and on activated T and B cells. PD-1 is thought to be involved in lymphocyte clonal selection and peripheral tolerance. The PD-1 ligands, PD-L1 (also known as B7-H1) and PD-L2 (B7-DC), are members ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [RMP1-14] |
| UniProt | Q02242 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA (order BSA/azide-free format for coating), Neutralization Studies (order BSA/azide-free format), Flow cytometry: 1-2ug/10 ⁶ cells in 0.1ml, Immunofluorescence: 0.5-1ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the PD1 antibody should be determined by the researcher. |