

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal NSJ-V9102-100UG**

Artikelname	Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal
Artikelnummer	NSJ-V9102-100UG
Hersteller Artikelnummer	V9102-100UG
Alternativnummer	NSJ-V9102-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 122-168 were used as the immunogen for this Bcl10 antibody.
Produktbeschreibung	Bcl10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1,14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of Bcl10...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM520]
UniProt	<a href="#">O95999</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)
Anwendungsbeschreibung	The optimal dilution of the Bcl10 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.