

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-CD68 Antibody, Clone: [SPM130], Mouse, Monoclonal NSJ-V9107IHC-7ML

Artikelname	Anti-CD68 Antibody, Clone: [SPM130], Mouse, Monoclonal
Artikelnummer	NSJ-V9107IHC-7ML
Hersteller Artikelnummer	V9107IHC-7ML
Alternativnummer	NSJ-V9107IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	A human alveolar macrophage subcellular fraction was used as the immunogen for this anti-CD68 antibody.
Produktbeschreibung	This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the r...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM130]
UniProt	P34810
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the anti-CD68 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.