

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Bromodeoxyuridine Antibody, Clone: [SPM166], Mouse, Monoclonal NSJ-V9115-100UG

Artikelname	Anti-Bromodeoxyuridine Antibody, Clone: [SPM166], Mouse, Monoclonal
Artikelnummer	NSJ-V9115-100UG
Hersteller Artikelnummer	V9115-100UG
Alternativnummer	NSJ-V9115-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	All
Immunogen	Bromodeoxyuridine was used as the immunogen for this anti-Bromodeoxyuridine antibody.
Produktbeschreibung	It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA sy...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM166]
UniProt	Not Applicable
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-Bromodeoxyuridine antibody for each application should be determined by the researcher.1. For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion with Trypsin at 1mg/ml PBS, 10 min at 37oC.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.