

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Plasma Cell Marker Antibody, Clone: [SPM310], Mouse, Monoclonal NSJ-V9116IHC-7ML

Artikelname	Plasma Cell Marker Antibody, Clone: [SPM310], Mouse, Monoclonal
Artikelnummer	NSJ-V9116IHC-7ML
Hersteller Artikelnummer	V9116IHC-7ML
Alternativnummer	NSJ-V9116IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Pancreatic cancer-related mucin was used as the immunogen for this Plasma Cell Marker antibody.
Produktbeschreibung	It recognizes an intra-cytoplasmic antigen, which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. Plasma cells, which are large lymphocytes derived from an antigen-specif...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM310]
UniProt	Not Known
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the Plasma Cell Marker antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. Not suitable for staining frozen tissues. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.