

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Plasma Cell Marker Antibody, Clone: [SPM310], Mouse, Monoclonal NSJ-V9116SAF-100UG

Artikelname	Plasma Cell Marker Antibody, Clone: [SPM310], Mouse, Monoclonal
Artikelnummer	NSJ-V9116SAF-100UG
Hersteller Artikelnummer	V9116SAF-100UG
Alternativnummer	NSJ-V9116SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Pancreatic cancer-related mucin was used as the immunogen for this Plasma Cell Marker antibody.
Produktbeschreibung	It recognizes an intra-cytoplasmic antigen, which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. Plasma cells, which are large lymphocytes derived from an antigen-specif...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM310]
UniProt	Not Known
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>The optimal dilution of the Plasma Cell Marker antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. Not suitable for staining frozen tissues.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>