

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 19 Antibody, Clone: [SPM561], Mouse, Monoclonal NSJ-V9127SAF-100UG

Artikelname	Cytokeratin 19 Antibody, Clone: [SPM561], Mouse, Monoclonal
Artikelnummer	NSJ-V9127SAF-100UG
Hersteller Artikelnummer	V9127SAF-100UG
Alternativnummer	NSJ-V9127SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Human mammary epithelial organoids were used as the immunogen for this Cytokeratin 19 antibody.
Produktbeschreibung	Cytokeratin-19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial m...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM561]
UniProt	<a href="#">P08727</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml,Flow cytometry: 1-2ug/10 <sup>6</sup> cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Cytokeratin 19 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.