

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Mouse IGFBP-7 (C-6His) EBT-EPT289

Artikelname	Recombinant Mouse IGFBP-7 (C-6His)
Artikelnummer	EBT-EPT289
Hersteller Artikelnummer	EPT289
Alternativnummer	EBT-EPT289-50
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Mouse Insulin-like Growth Factor-binding Protein 7 is produced by our Mammalian expression system and the target gene encoding Ser26-Leu281 is expressed with a 6His tag at the C-terminus....
Molekulargewicht	Molecular weight: 27.2 KDa. Apparent molecular weight: 32-35 KDa, reducing conditions
UniProt	Q61581
Reinheit	Greater than 95% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Background: Insulin-like growth factor-binding protein 7 (IGFBP-7) is a secreted glycosylated protein that contains three protein domain modules. IGFBP7 contains an N-terminal IGFBP domain, followed by a Kazal-type serine proteinase inhibitor domain and a C-terminal immunoglobulin-like C2-type domain. Human and mouse IGFBP7 are highly homologous and share 94% aa sequence identity. It is expressed in many normal tissues and in cancer cells. It is abundantly expressed in high endothelial venules (HEVs) of blood vessels in the secondary lymphoid tissues. It binds IGF and insulin with very low affinity and has been shown to enhance the mitogenic actions of IGF and insulin. IGFBP7 also has IGF/insulin-independent activities. It interacts with heparan sulfate proteoglycans, type IV collagen, and specific chemokines. It supports weak cell adhesion, promotes cell spreading on type IV collagen, and stimulates the production of the potent vasodilator PGI₂. It modulates tumor cell growth and has also been implicated in angiogenesis.