

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mesothelin Antibody, IgG2a, Clone: [MN-1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344444

Artikelname	Mesothelin Antibody, IgG2a, Clone: [MN-1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage
Artikelnummer	BYT-ORB344444
Hersteller Artikelnummer	orb344444
Alternativnummer	BYT-ORB344444-100
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FC, IHC, WB
Spezies Reaktivität	Human
Immunogen	This antibody was produced in mesothelin-deficient mice by immunizations with plasmid cDNA encoding human MSLN full length protein followed by a single boost of a recombinant human mesothelin-Fc fusion protein.
Konjugation	Unconjugated
Produktbeschreibung	Mesothelin antibody...
Klonalität	Monoclonal
Konzentration	1.0 mg/mL
Klon-Bezeichnung	[MN-1]

Isotyp	IgG2a
NCBI	005814
Pubmed	https://pubmed.ncbi.nlm.nih.gov/27544053 target=_blank rel=nofollow>27544053
UniProt	Q13421
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This antibody is directed against human mesothelin protein. This product was purified from tissue culture supernatant fluid by Protein A chromatography. Cross reactivity with homologues from other sources has not been tested.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:10,000 - 1:50,000, FC: 1:800, IHC: 1:100, WB: 1:1,000
Anwendungsbeschreibung	Application Notes: This antibody has been tested for use in immunohistochemistry, flow cytometry, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 40 kDa in size corresponding to mature mesothelin by western blotting in the appropriate cell lysate or extract. For immunohistochemistry, archival PEFF human tissues were deparaffinized followed by hydration. Antigen-retrieval is recommended. Block tissues with 1% BSA in PBS for 30 min at 23 C. Antibodies are diluted in 1% BSA and reacted with tissue for 60 min at room temperature