

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mesothelin Antibody, IgG2a, Clone: [MN-1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344445

| | |
|--------------------------|--|
| Artikelname | Mesothelin Antibody, IgG2a, Clone: [MN-1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage |
| Artikelnummer | BYT-ORB344445 |
| Hersteller Artikelnummer | orb344445 |
| Alternativnummer | BYT-ORB344445-25 |
| Hersteller | Biorbyt |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IHC, IP, WB |
| Spezies Reaktivität | Human |
| Immunogen | This antibody was produced in mesothelin-deficient mice by immunizations with plasmid cDNA encoding human MSLN full length protein followed by a single boost of a recombinant human mesothelin-Fc fusion protein. |
| Konjugation | Unconjugated |
| Produktbeschreibung | Mesothelin antibody... |
| Klonalität | Monoclonal |
| Konzentration | 1.0 mg/mL |
| Klon-Bezeichnung | [MN-1] |

| | |
|------------------------|--|
| Isotyp | IgG2a |
| NCBI | 53988378 |
| UniProt | Q13421 |
| Puffer | Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This antibody is directed against human mesothelin protein. This product was purified from tissue culture supernatant fluid by Protein A chromatography. Cross reactivity with homologues from other sources has not been tested. |
| Formulierung | Liquid (sterile filtered) |
| Application Verdünnung | ELISA: 1:10,000 - 1:50,000, IHC: 1:100, IP: 1:800, WB: 1:1,000 |
| Anwendungsbeschreibung | Application Notes: This antibody has been tested for use in immunohistochemistry, flow cytometry, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 40 kDa in size corresponding to mature mesothelin by western blotting in the appropriate cell lysate or extract. For immunohistochemistry, archival PEFF human tissues were deparaffinized followed by hydration. Antigen-retrieval is recommended. Block tissues with 1% BSA in PBS for 30 min at 23 C. Antibodies are diluted in 1% BSA and reacted with tissue for 60 min at room temperature |