

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AKT phospho S473 Antibody, IgG1, Clone: [17F6.B11], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344456

Artikelname	AKT phospho S473 Antibody, IgG1, Clone: [17F6.B11], Unconjugated, Mouse, Monoclonal Preis auf Anfrage
Artikelnummer	BYT-ORB344456
Hersteller Artikelnummer	orb344456
Alternativnummer	BYT-ORB344456-1
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IF, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	This monoclonal antibody was produced by repeated immunizations with a synthetic peptide corresponding to residues surrounding S473 of human AKT1 protein.
Konjugation	Unconjugated
Produktbeschreibung	Akt (phospho-S473) antibody...
Klonalität	Monoclonal
Konzentration	1.0 mg/mL
Klon-Bezeichnung	[17F6.B11]

Isotyp	IgG1
NCBI	62241011
UniProt	P31749
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was purified from concentrated tissue culture supernate by Protein A chromatography. This antibody is specific for human and mouse AKT protein phosphorylated at S473. A BLAST analysis was used to suggest cross-reactivity with AKT pS473 from human, mouse, rat and chimpanzee sources based on 100% homology with the immunizing sequence. Cross-reactivity with AKT from other sources has not been determined. Cross-reactivity with AKT2 and AKT3 has not been determined.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:20,000, IHC: 20 µg/ml, IF: 1:500 - 1:3,000, WB: 1:500 - 1:3,000
Anwendungsbeschreibung	Application Notes: This monoclonal antibody is tested in ELISA, immunohistochemistry, immunofluorescent microscopy, and western blotting. Expect a band approximately 56 kDa in size corresponding to phosphorylated AKT protein by western blotting in the appropriate cell lysate or extract. This phospho-specific monoclonal antibody reacts with human and mouse AKT pS473 and shows minimal reactivity by ELISA against the non-phosphorylated form of the immunizing peptide. Specific conditions for reactivity should be optimized by the end user. For immunohistochemistry use formalin-fixed paraffin-embedded sections. No pre-treatment of sample is required. Cell Signaling, Cancer, Neuroscience, Signal Transduction research