

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Alpha-Tubulin Antibody, Unconjugated, Rabbit, Polyclonal Preis auf Anfrage BYT-ORB345510

Artikelname	Alpha-Tubulin Antibody, Unconjugated, Rabbit, Polyclonal Preis auf Anfrage
Artikelnummer	BYT-ORB345510
Hersteller Artikelnummer	orb345510
Alternativnummer	BYT-ORB345510-200
Hersteller	Biorbyt
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Anti-Tubulin Loading Control Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the C-Terminal region near amino acids 425-451 of Human alpha Tubulin.
Konjugation	Unconjugated
Produktbeschreibung	Alpha-Tubulin antibody...
Klonalität	Polyclonal
Konzentration	1.1 mg/mL
NCBI	17986283

UniProt	P68363
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Tubulin Loading Control Antibody is directed against human alpha Tubulin protein. The Loading Control Antibody was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest that this antibody would react with alpha Tubulin from a wide range of organisms, including avian, mammalian aquatic, parasitic and alga sources based on 100% homology for the immunogen sequence. Cross reactivity will occur with all isoforms of alpha tubulin. Such broad reactivity makes this antibody useful as an excellent loading control.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:90,000, IHC: 1:500 - 1:2,000, IF: 1:500 - 1:2,000, WB: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Application Notes: Anti-Tubulin Antibody has been tested for use in ELISA, immunofluorescence, and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band at ~50 kDa in size corresponding to alpha tubulin by western blotting in most cell lysates or extracts