

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Bovine IgG F(ab)2 Antibody Fluorescein Conjugated, FITC, Goat, Polyclonal BYT-ORB346550

Artikelname	Bovine IgG F(ab)2 Antibody Fluorescein Conjugated, FITC, Goat, Polyclonal
Artikelnummer	BYT-ORB346550
Hersteller Artikelnummer	orb346550
Alternativnummer	BYT-ORB346550-20
Hersteller	Biorbyt
Wirt	Goat
Kategorie	Antikörper
Applikation	FC, FLISA, IF
Spezies Reaktivität	Bovine
Immunogen	Bovine IgG F(ab)2 fragment
Konjugation	FITC
Produktbeschreibung	Bovine IgG F(ab)2 antibody (FITC)...
Klonalität	Polyclonal
Puffer	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Reinheit	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Goat Serum, Bovine IgG, Bovine IgG F(ab)2 and Bovine Serum. No reaction was observed against Bovine IgG F(c).
Formulierung	Lyophilized
Application Verdünnung	FLISA: 1:10,000 - 1:1:50,000, FC: 1:500 - 1:2,500, IF: 1:1,000 - 1:1:5,000
Anwendungsbeschreibung	<p>Application Notes: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1×10^6 cells in flow cytometry is approximately 1.0 μg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications</p>