

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AKT phospho S473 Antibody, Unconjugated, Rabbit, Polyclonal Preis auf Anfrage BYT-ORB420307

Artikelname	AKT phospho S473 Antibody, Unconjugated, Rabbit, Polyclonal Preis auf Anfrage
Artikelnummer	BYT-ORB420307
Hersteller Artikelnummer	orb420307
Alternativnummer	BYT-ORB420307-25
Hersteller	Biorbyt
Wirt	Rabbit
Kategorie	Antikörper
Applikation	DOT, ELISA, IF, IHC, Multiplex Assay, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Rabbit Anti-AKTpS473 Antibody was prepared by repeated immunizations in rabbits with a synthetic peptide corresponding to a C-terminus region near phospho Serine 473 of the human, mouse, rat and chicken AKT proteins conjugated to KLH.
Konjugation	Unconjugated
Produktbeschreibung	AKT pS473 antibody...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
NCBI	62241011

UniProt	P31749
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using a phospho peptide coupled to agarose beads followed by solid phase adsorption(s) against a non-phospho peptide and non-specific peptide to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. This antibody is specific for phosphorylated human AKTpS473. Minimal reactivity occurs against non-phosphorylated AKT. Reactivity against AKT from other species may occur but has not yet been tested.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:15,000 - 1:60,000, IHC: 1:100 - 1:500, WB: 1:200 - 1:1000
Anwendungsbeschreibung	Application Notes: This AKT antibody is phospho specific for pS473 and is tested in ELISA, western blotting, immunohistochemistry (formalin-fixed paraffin-embedded sections), and immunofluorescence. By immunoblot a single band of the expected apparent molecular weight ~56kDa is observed. For immunohistochemistry no pre-treatment of sample is required