

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27Kip1 Antibody, Clone: [KIP1/769], Unconjugated, Mouse, Monoclonal BYT-ORB2638253

| | |
|--------------------------|---|
| Artikelname | p27Kip1 Antibody, Clone: [KIP1/769], Unconjugated, Mouse, Monoclonal |
| Artikelnummer | BYT-ORB2638253 |
| Hersteller Artikelnummer | orb2638253 |
| Alternativnummer | BYT-ORB2638253-100 |
| Hersteller | Biorbyt |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant human protein was used as the immunogen for the p27Kip1 antibody. |
| Konjugation | Unconjugated |
| Produktbeschreibung | This mAb recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27Kip1 functions as a negative regulator of G1 progre... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KIP1/769] |
| UniProt | P46527 |

| | |
|------------------------|--|
| Puffer | 1 mg/ml in 1X PBS, rAlbumin free, sodium azide free |
| Application Verdünnung | Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Application Notes: Optimal dilution of the p27Kip1 antibody to be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min |