

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal BYT-ORB2638903

| | |
|--------------------------|---|
| Artikelname | MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal |
| Artikelnummer | BYT-ORB2638903 |
| Hersteller Artikelnummer | orb2638903 |
| Alternativnummer | BYT-ORB2638903-100 |
| Hersteller | Biorbyt |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Human milk-fat globule membranes (HMFGM) was used as the immunogen for the MUC1 antibody. |
| Konjugation | Unconjugated |
| Produktbeschreibung | In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of MUC1/Mucin-1/Epithelial Marker Antigen/EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhes... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MUC1/520] |

| | |
|------------------------|--|
| UniProt | P15941 |
| Puffer | 1 mg/ml in 1X PBS, rAlbumin free, sodium azide free |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Western blot: 1-2ug/ml |
| Anwendungsbeschreibung | Application Notes: Optimal dilution of the MUC1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min |