

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant EpCAM Antibody / Extracellular domain, Clone: [rEGP40/1372], Unconjugated, Mouse BYT-ORB533882

| | |
|--------------------------|---|
| Artikelname | Recombinant EpCAM Antibody / Extracellular domain, Clone: [rEGP40/1372], Unconjugated, Mouse |
| Artikelnummer | BYT-ORB533882 |
| Hersteller Artikelnummer | orb533882 |
| Alternativnummer | BYT-ORB533882-20,BYT-ORB533882-100 |
| Hersteller | Biorbyt |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Canine, Feline, Human |
| Immunogen | A portion of amino acids 100-224 (extracellular domain) was used as the immunogen for the recombinant EpCAM antibody. |
| Konjugation | Unconjugated |
| Produktbeschreibung | EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast major... |
| Klonalität | Recombinant |
| Klon-Bezeichnung | [rEGP40/1372] |

| | |
|------------------------|--|
| UniProt | P16422 |
| Puffer | 0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide |
| Application Verdünnung | Western blot: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Application Notes: Optimal dilution of the recombinant EpCAM antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min |