

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **H1F0 Antibody, Clone: [1415-1], Unconjugated, Mouse, Monoclonal BYT-ORB1252534**

Artikelname	H1F0 Antibody, Clone: [1415-1], Unconjugated, Mouse, Monoclonal
Artikelnummer	BYT-ORB1252534
Hersteller Artikelnummer	orb1252534
Alternativnummer	BYT-ORB1252534-100
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC-P, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Nuclei of human leukemia biopsy cells were used as the immunogen for the Histone antibody.
Konjugation	Unconjugated
Produktbeschreibung	H1F0 Antibody...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[1415-1]
UniProt	<a href="#">P07305</a>
Puffer	PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide

Formulierung	Liquid
Target-Kategorie	H1F0
Anwendungsbeschreibung	<p>Application Notes: Western blot: 0.5-1 µg/mL Flow Cytometry: 0.5-1 µg/million cells in 0.1ml Immunofluorescence: 0.5-1 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 min at RT (1) Prediluted format : incubate for 30 min at RT (2) Optimal dilution of the Histone antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p>