

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD63 Antibody, Clone: [MX-49.129.5], Unconjugated, Mouse, Monoclonal BYT-ORB1252797

| | |
|--------------------------|--|
| Artikelname | CD63 Antibody, Clone: [MX-49.129.5], Unconjugated, Mouse, Monoclonal |
| Artikelnummer | BYT-ORB1252797 |
| Hersteller Artikelnummer | orb1252797 |
| Alternativnummer | BYT-ORB1252797-100 |
| Hersteller | Biorbyt |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Full length human CD63 was used as the immunogen for this antibody. |
| Konjugation | Unconjugated |
| Produktbeschreibung | CD63 Antibody... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [MX-49.129.5] |
| UniProt | P08962 |
| Puffer | PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide |

| | |
|------------------------|---|
| Formulierung | Liquid |
| Target-Kategorie | CD63 |
| Anwendungsbeschreibung | <p>Application Notes: Flow Cytometry: 0.5-1 ug/million cells IF: 0.5-1 µg/mL WB: 1-2 µg/mL IHC (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2) The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titrated up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p> |