

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TIMP1 Antibody, Unconjugated, Rabbit, Polyclonal BYT-ORB1272681**

Artikelname	TIMP1 Antibody, Unconjugated, Rabbit, Polyclonal
Artikelnummer	BYT-ORB1272681
Hersteller Artikelnummer	orb1272681
Alternativnummer	BYT-ORB1272681-0.1
Hersteller	Biorbyt
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, NeA, WB
Spezies Reaktivität	Human
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hTIMP-1 (human Tissue Inhibitor of Metalloproteinases-1). Human TIMP-1 specific antibody was purified by affinity chromatography employing immobilized hTIMP-1 matrix.
Konjugation	Unconjugated
Produktbeschreibung	TIMP1 Antibody...
Klonalität	Polyclonal
Konzentration	batch dependent
NCBI	<a href="#">01033</a>
UniProt	<a href="#">P01033</a>

Formulierung	Lyophilized
Target-Kategorie	TIMP1
Anwendungsbeschreibung	<p>Application Notes: Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Human TIMP-1 (3.0 µg/mL), a concentration of 5.0 µg/mL of this antibody is required. ELISA: Indirect: To detect hTIMP-1 by indirect ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTIMP-1. Sandwich To detect hTIMP-1 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our biotinylated Anti-Human TIMP-1 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTIMP-1. Western Blot: To detect hTIMP-1 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTIMP-1 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions</p>