

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27Kip1 Antibody, Clone: [SX53G8], Unconjugated, Mouse, Monoclonal BYT-ORB248293

Artikelname	p27Kip1 Antibody, Clone: [SX53G8], Unconjugated, Mouse, Monoclonal
Artikelnummer	BYT-ORB248293
Hersteller Artikelnummer	orb248293
Alternativnummer	BYT-ORB248293-20,BYT-ORB248293-100
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Purified GST-p27 fusion protein of human origin was used as the immunogen for this p27Kip1 antibody.
Konjugation	Unconjugated
Produktbeschreibung	This antibody recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. The p27Kip1 antibody is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27Kip1 functions as a negative...
Klonalität	Monoclonal
Klon-Bezeichnung	[SX53G8]

Puffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Application Verdünnung	Flow cytometry: 0.5-1ug/million cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>Application Notes: The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the p27Kip1 antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p>