

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Monoclonal Antibody to Bromodeoxyuridine (BrdU) (Proliferation Marker)(Clone : BRD.3), Clone: [BRD.3], Mouse ABI-36-11106**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Monoclonal Antibody to Bromodeoxyuridine (BrdU) (Proliferation Marker)(Clone : BRD.3), Clone: [BRD.3], Mouse                                                                                                                                                  |
| Artikelnummer            | ABI-36-11106                                                                                                                                                                                                                                                  |
| Hersteller Artikelnummer | 36-11106                                                                                                                                                                                                                                                      |
| Alternativnummer         | ABI-36-11106-100UG                                                                                                                                                                                                                                            |
| Hersteller               | Abeomics                                                                                                                                                                                                                                                      |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FACS, IF, IHC                                                                                                                                                                                                                                                 |
| Spezies Reaktivität      | All                                                                                                                                                                                                                                                           |
| Immunogen                | Bromodeoxyuridine (BrdU) conjugated to BSA                                                                                                                                                                                                                    |
| Produktbeschreibung      | It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA sy... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [BRD.3]                                                                                                                                                                                                                                                       |
| Reinheit                 | Affinity Chromatography                                                                                                                                                                                                                                       |
| Formulierung             | Purified                                                                                                                                                                                                                                                      |

|                        |                                                                                                                                                                                                                                                                |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target-Kategorie       | Bromodeoxyuridine (BrdU) (Proliferation Marker)                                                                                                                                                                                                                |
| Application Verdünnung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)[For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion with |