

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Anti-Complement 4d (C4d) (Acute Humoral Rejection Marker) Monoclonal Antibody(Clone: C4D204)-CF488, Clone: [C4D204], Mouse ABI-36-3314**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Anti-Complement 4d (C4d) (Acute Humoral Rejection Marker) Monoclonal Antibody(Clone: C4D204)-CF488, Clone: [C4D204], Mouse                                                                                                                                    |
| Artikelnummer            | ABI-36-3314                                                                                                                                                                                                                                                   |
| Hersteller Artikelnummer | 36-3314                                                                                                                                                                                                                                                       |
| Alternativnummer         | ABI-36-3314-0.5AT100MLATUG/ML                                                                                                                                                                                                                                 |
| Hersteller               | Abeomics                                                                                                                                                                                                                                                      |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | ELISA, IF, IHC                                                                                                                                                                                                                                                |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human Complement 4d protein                                                                                                                                                                                                                       |
| Konjugation              | CF488                                                                                                                                                                                                                                                         |
| Produktbeschreibung      | This MAb is specific to Complement 4d (C4d) and reacts with the secreted as well as cell-bound C4d.C4d is a degradation product of the activated complement factor C4b. Complement 4b is typically activated by binding of Abs to specific target molecule... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [C4D204]                                                                                                                                                                                                                                                      |

|                        |                                                                                                                                                                                                                                                                 |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NCBI                   | <a href="#">720</a>                                                                                                                                                                                                                                             |
| UniProt                | P00L4, P00L5                                                                                                                                                                                                                                                    |
| Application Verdünnung | ELISA (For coating use Ab at 1-2ug/ml order Ab without BSA), Immunofluorescence (1:50-1:100),Immunohistochemistry (Formalin-fixed) (1:200-1:400 for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8 |