

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Swine anti Goat IgA (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAG/IGA(FC)

Artikelname	Swine anti Goat IgA (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal
Artikelnummer	NMB-SWAG/IGA(FC)
Hersteller Artikelnummer	SwAG/IgA(Fc)
Alternativnummer	NMB-SWAG/IGA(FC)
Hersteller	NordicMubio
Wirt	Porcine
Kategorie	Antikörper
Spezies Reaktivität	Goat
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgA
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is restricted to the Fc part of the IgA molecule as tested in immunoelectrophoresis and radial immunodiffusion. In immunoelectrophoresis using various antiserum concentrations against normal goat serum only IgA is prec...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized
Quelle	Highly purified secretory IgA isolated from pooled goat milk. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter centre well and 2 μ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.