

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu CD3 zeta (pY72) Purified, Clone: [EM-26], Monoclonal EXB-11-712-C100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu CD3 zeta (pY72) Purified, Clone: [EM-26], Monoclonal |
| Artikelnummer | EXB-11-712-C100 |
| Hersteller Artikelnummer | 11-712-C100 |
| Alternativnummer | EXB-11-712-C100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC, ICC, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | A phospho specific peptide corresponding to the amino acids surrounding tyrosine 72 of mouse CD3 zeta linked to KLH |
| Produktbeschreibung | CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |
| Klon-Bezeichnung | [EM-26] |
| Isotyp | Mouse IgG2b |
| Puffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|--|
| Lagerung | 2°C to 8°C |
| Target-Kategorie | CD3 zeta (pY72) |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western blotting: Recommended dilution: 2 - 5 µg/ml, positive control: Jurkat cells lysate treated with pervanadate, splenocyte lysate of Balb/c or F1 mouse treated with pervanadate, non-reducing conditions recommended. Flow cytometry: Intracellular staining, recommended dilution: 1-9 µg/ml, positive control: Jurkat cells treated with pervanadate, T-cells from lymph nodes of OT-1 mouse treated with pervanadate. |