

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse IgG1 Isotype Control APC, Clone: [MOPC-21], Monoclonal EXB-1A-632-C100

| | |
|--------------------------|---|
| Artikelname | Mouse IgG1 Isotype Control APC, Clone: [MOPC-21], Monoclonal |
| Artikelnummer | EXB-1A-632-C100 |
| Hersteller Artikelnummer | 1A-632-C100 |
| Alternativnummer | EXB-1A-632-C100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Konjugation | APC |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/ml |
| Klon-Bezeichnung | [MOPC-21] |
| Isotyp | Mouse IgG1 kappa |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Lagerung | 2°C to 8°C |
| Antibody Type | Monoclonal Antibody |

| | |
|------------------------|---|
| Application Verdünnung | Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the |
| Anwendungsbeschreibung | Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg/ml of incubation mixture), change the conditions of your experiment to reduce the background. |