

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit F(ab)2 anti-Goat IgG (H+L)-unconj., MinX none DNA-SEC-182666

|                                  |                                                                                                                                                                                                                                                               |
|----------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname                      | Rabbit F(ab)2 anti-Goat IgG (H+L)-unconj., MinX none                                                                                                                                                                                                          |
| Artikelnummer                    | DNA-SEC-182666                                                                                                                                                                                                                                                |
| Hersteller Artikelnummer         | SEC-182666                                                                                                                                                                                                                                                    |
| Alternativnummer                 | DNA-SEC-182666                                                                                                                                                                                                                                                |
| Hersteller                       | dianova                                                                                                                                                                                                                                                       |
| Wirt                             | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                        | Antikörper                                                                                                                                                                                                                                                    |
| Applikation                      | ELISA,IHC,WB                                                                                                                                                                                                                                                  |
| Spezies Reaktivität              | Goat                                                                                                                                                                                                                                                          |
| Immunogen                        | Goat IgG whole molecule                                                                                                                                                                                                                                       |
| Konjugation                      | Unconjugated                                                                                                                                                                                                                                                  |
| Format                           | F(ab')2                                                                                                                                                                                                                                                       |
| Spezifität                       | IgG (H+L)                                                                                                                                                                                                                                                     |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion                                                                                                                                                                                                                                           |
| Produktbeschreibung              | F(ab)2 Anti-Goat IgG Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techni... |
| Klonalität                       | Polyclonal                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Konzentration          | 10.0 mg/mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Isotyp                 | Ig                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Puffer                 | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Reinheit               | This product is a F(ab') <sub>2</sub> fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.                                                                                          |
| Formulierung           | Lyophilized                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Formel                 | 10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN <sub>3</sub>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Target-Kategorie       | Goat                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Antibody Type          | Secondary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |