

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none DNA-SEC-182668

Artikelname	Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none
Artikelnummer	DNA-SEC-182668
Hersteller Artikelnummer	SEC-182668
Alternativnummer	DNA-SEC-182668
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FLISA,FACS,IF
Spezies Reaktivität	Goat
Immunogen	Goat IgG whole molecule
Konjugation	FITC
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Anti-Goat IgG Fluorescein Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunoche...

Klonalität	Polyclonal
Konzentration	10.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product is a F(ab') ₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.
Formulierung	Lyophilized
Formel	10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN ₃
Target-Kategorie	Goat
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500-1:2,500, Fluorochrome Protein Value: 4.0, IF Microscopy Dilution: 1:1,000-1:5,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.