

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (H+L)-HRPO, MinX none DNA-SEC-182706

Artikelname	Goat F(ab)2 anti-Rat IgG (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-182706
Hersteller Artikelnummer	SEC-182706
Alternativnummer	DNA-SEC-182706
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rat
Immunogen	Rat IgG whole molecule
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper...

Klonalität	Polyclonal
Konzentration	10.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product is a F(ab') ₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Goat IgG F(c) or anti-Pepsin.
Formulierung	Lyophilized
Formel	10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target-Kategorie	Rat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	F(ab) ₂ Anti-Rat IgG (H&L) Peroxidase secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.