

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Goat IgG (Fc)-HRPO, MinX none DNA-SEC-182852

Artikelname	Rabbit IgG anti-Goat IgG (Fc)-HRPO, MinX none
Artikelnummer	DNA-SEC-182852
Hersteller Artikelnummer	SEC-182852
Alternativnummer	DNA-SEC-182852
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Goat
Immunogen	Anti-Goat IgG F(c) was produced by repeated immunization with Goat IgG F(c) fragment in rabbit.
Konjugation	HRPO
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Goat IgG F(ab)2 Peroxidase Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 molecules lack the F...

Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Goat IgG, Goat IgG F(c) and Goat Serum. No reaction was observed against Goat IgG F(ab) .
Formulierung	Lyophilized
Formel	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Goat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:30,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	Goat IgG F(c) peroxidase conjugated antibody has been assayed against 1.0 ug of Goat IgG in a standard capture ELISA using TMB as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:16,000 is suggested for this product.