

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Guinea Pig IgG (H+L)-HRPO, MinX none DNA-SEC-182923

Artikelname	Rabbit IgG anti-Guinea Pig IgG (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-182923
Hersteller Artikelnummer	SEC-182923
Alternativnummer	DNA-SEC-182923
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Guinea pig
Immunogen	Anti-Guinea Pig IgG was produced by repeated immunization with guinea pig IgG whole molecule in rabbit.
Konjugation	HRPO
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Guinea Pig IgG Peroxidase Antibody generated in rabbit detects guinea pig IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bact...

Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Guinea Pig IgG and Guinea Pig Serum.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Guinea Pig
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:500 - 2,500, Western Blot Dilution: 1:2,000 - 10,000
Anwendungsbeschreibung	Antibody Anti-Guinea Pig IgG (H&L) peroxidase conjugated has been assayed against 1.0 ug of Guinea Pig IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.