

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Horse IgG (Fc)-Biotin, MinX none DNA-SEC-182985

Artikelname	Rabbit IgG anti-Horse IgG (Fc)-Biotin, MinX none
Artikelnummer	DNA-SEC-182985
Hersteller Artikelnummer	SEC-182985
Alternativnummer	DNA-SEC-182985
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Equine
Immunogen	Horse IgG F(c) fragment
Konjugation	Biotin
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Horse IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of horse IgG and o...
Klonalität	Polyclonal

Konzentration	1.5 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Horse IgG, Horse IgG F(c) and Horse Serum. No reaction was observed against Horse IgG F(ab).
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Horse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:15,000 of the reconstitution concentration is suggested for this product.