

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Human Kappa light chain-unconj., MinX none DNA-SEC-183008

Artikelname	Goat IgG anti-Human Kappa light chain-unconj., MinX none
Artikelnummer	DNA-SEC-183008
Hersteller Artikelnummer	SEC-183008
Alternativnummer	DNA-SEC-183008
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human kappa light chain
Konjugation	Unconjugated
Format	IgG
Spezifität	Kappa (light chain)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The anti-Human kappa (kappa chain) Antibody detects the kappa chain subunit. Immunoglobulins are heterotetramers composed of 2 immunoglobulin heavy and 2 immunoglobulin light chains. The immunoglobulin light chain is the small polypeptide subunit of ...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0
Reinheit	Human κ (kappa chain) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. Specificity for kappa chain was confirmed by ELISA minimal cross reactivity against other human heavy or lambda light chain isotypes.
Formulierung	Liquid (sterile filtered)
Formel	125 mM Sodium Borate, 75 mM NaCl, 5 mM EDTA, pH 8.0, sterile filtered, 0.01% NaN ₃
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:20,000, Immunohistochemistry Dilution: 1:500 - 1:2,000, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	Human κ (kappa chain) Antibody is suitable for immunoassays where specificity to the immunoglobulin kappa subunit is desired. Antibody has been tested by ELISA, western blot immunoblot, and immunohistochemistry. Optimal concentrations in immunoassays should be determined by the researcher.