

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Human IgG (F(ab)2)-Alk. Phos., MinX Ms DNA-SEC-183084

Artikelname	Rabbit IgG anti-Human IgG (F(ab)2)-Alk. Phos., MinX Ms
Artikelnummer	DNA-SEC-183084
Hersteller Artikelnummer	SEC-183084
Alternativnummer	DNA-SEC-183084
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgG F(ab)2 fragment
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	Mouse
Produktbeschreibung	Anti-Human IgG F(ab)2 Alkaline Phosphatase Antibody generated in rabbit recognizes the dimeric Fab portion of the human IgG molecule. Human IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme p...

Klonalität	Polyclonal
Konzentration	1.4 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Human IgG, Human IgG F(ab') ₂ and Human Serum. No reaction was observed against Human IgG F(c) or Mouse Serum Proteins.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl, 0.1 mM ZnCl, 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.01% NaN ₃
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 µg of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,500 to 1:11,000 of the stated concentration is suggested for this product.