

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Human IgA-Biotin, MinX none DNA-SEC-183089

Artikelname	Rabbit IgG anti-Human IgA-Biotin, MinX none
Artikelnummer	DNA-SEC-183089
Hersteller Artikelnummer	SEC-183089
Alternativnummer	DNA-SEC-183089
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgA alpha heavy chain
Konjugation	Biotin
Format	IgG
Spezifität	IgA
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Human IgA Biotin Antibody generated in rabbit detects immunoglobulin A (alpha chain) from human. Immunoglobulin A (IgA) is an antibody that plays a critical role in mucosal immunity. IgA has two subclasses (IgA1 and IgA2) and can exist in a dime...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Human IgA and Human Serum. No reaction was observed against Human IgG or IgM. Specificity was confirmed by ELISA minimal cross reactivity against other human heavy or light chain isotypes.
Formulierung	Lyophilized
Formel	10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Thimerosal
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:5,000 - 1:20,000, Immunohistochemistry Dilution: 1:250 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Human IgA in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:100,000 of the reconstitution concentration is suggested for this product.