

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rabbit IgG (Fc)-Biotin, MinX none DNA-SEC-183838

Artikelname	Goat F(ab)2 anti-Rabbit IgG (Fc)-Biotin, MinX none
Artikelnummer	DNA-SEC-183838
Hersteller Artikelnummer	SEC-183838
Alternativnummer	DNA-SEC-183838
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Rabbit IgG F(c) fragment
Konjugation	Biotin
Format	F(ab')2
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Anti-Rabbit IgG F(c) Biotin Antibody was generated in goat and detects specifically Rabbit IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	F(ab') fragment biotin conjugated secondary antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Rabbit IgG F(ab).
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:12,000 to 1:60,000 of the reconstitution concentration is suggested for this product. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.