

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey F(ab)2 anti-Rabbit IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh DNA-SEC-183844

Artikelname	Donkey F(ab)2 anti-Rabbit IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh
Artikelnummer	DNA-SEC-183844
Hersteller Artikelnummer	SEC-183844
Alternativnummer	DNA-SEC-183844
Hersteller	dianova
Wirt	Donkey
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Rabbit IgG whole molecule
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rat,Sheep

Produktbeschreibung	F(ab)2 Anti-Rabbit IgG (H&L) Antibody generated in donkey detects immunoglobulin g from Rabbit, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunogl...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:8,000 - 1:40,000, Immunohistochemistry Dilution: 1:500 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	F(ab)2 Anti-Rabbit IgG Peroxidase Conjugated Secondary Antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, as well as other antibody detection methods.