

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (Fc)-HRPO, MinX Bo,Ho,Hu DNA-SEC-183861

Artikelname	Goat F(ab)2 anti-Rat IgG (Fc)-HRPO, MinX Bo,Ho,Hu
Artikelnummer	DNA-SEC-183861
Hersteller Artikelnummer	SEC-183861
Alternativnummer	DNA-SEC-183861
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rat
Immunogen	Rat IgG F(c) fragment
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	Bovine,Equine,Human
Produktbeschreibung	F(ab)2 Anti-Rat IgG F(c) Peroxidase Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must b...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab) or Bovine, Horse, Human Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Rat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:50,000 - 1:100,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:200,000 of the reconstitution concentration is suggested for this product for ELISA and Western blots and a 1:500 to 1:5,000 for immunohistochemistry on tissue sections.