

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (Fc)-Alk. Phos., MinX Bo,Ho,Hu DNA-SEC-183864

Artikelname	Goat F(ab)2 anti-Rat IgG (Fc)-Alk. Phos., MinX Bo,Ho,Hu
Artikelnummer	DNA-SEC-183864
Hersteller Artikelnummer	SEC-183864
Alternativnummer	DNA-SEC-183864
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rat
Immunogen	Rat IgG F(c) fragment
Konjugation	Alk. Phos.
Format	F(ab')2
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	Bovine,Equine,Human
Produktbeschreibung	F(ab)2 Anti-Rat IgG F(c) Alkaline Phosphatase Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, considerat...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	F(ab') ₂ Anti-Rat IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Rat IgG, Rat IgG Fc and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat F(ab) or Bovine, Horse, and Human Serum Proteins.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl, 0.1 mM ZnCl, 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.05% NaN ₃
Target-Kategorie	Rat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	F(ab') ₂ Anti-Rat IgG Antibody has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for F(ab') ₂ Anti-Rat IgG Antibody.