

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Sheep IgG (H+L)-unconj., MinX none DNA-SEC-183892

Artikelname	Rabbit F(ab)2 anti-Sheep IgG (H+L)-unconj., MinX none
Artikelnummer	DNA-SEC-183892
Hersteller Artikelnummer	SEC-183892
Alternativnummer	DNA-SEC-183892
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Sheep
Immunogen	Sheep IgG whole molecule
Konjugation	Unconjugated
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper...

Klonalität	Polyclonal
Konzentration	1 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Sheep IgG and Sheep Serum. No reaction was observed against anti-Pepsin and anti-Rabbit IgG F(c).
Formulierung	Liquid (sterile filtered)
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target-Kategorie	Sheep
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000-1:5,000, Western Blot Dilution: 1:2,000-1:10,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1×10^6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.