

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Sheep IgG (H+L)-Alk. Phos., MinX none DNA-SEC-183895

Artikelname	Rabbit F(ab)2 anti-Sheep IgG (H+L)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183895
Hersteller Artikelnummer	SEC-183895
Alternativnummer	DNA-SEC-183895
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Sheep
Immunogen	Sheep IgG whole molecule
Konjugation	Alk. Phos.
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency....
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Rabbit Serum, Sheep IgG and Sheep Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Sheep
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 µg of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.