

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **COMBI IC Reagent: Mouse anti CD45 (FITC) and Mouse anti CD14 (PE), Clone: [VIT200 and MEM18], FITC/PE, Monoclonal NMB-GCT-201**

Article Name	COMBI IC Reagent: Mouse anti CD45 (FITC) and Mouse anti CD14 (PE), Clone: [VIT200 and MEM18], FITC/PE, Monoclonal
Biozol Catalog Number	NMB-GCT-201
Supplier Catalog Number	GCT-201
Alternative Catalog Number	NMB-GCT-201
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	FC, IF
Species Reactivity	Human
Conjugation	FITC/PE
Product Description	The CD45 molecule is typically expressed at high levels on all hematopoietic cells. CD45 is a major component of the glycocalix of these cells and can be expressed in different isoforms. Antibody VIT200 recognizes a pan CD45 epitope, which is express...
Clonality	Monoclonal
Clone Designation	[VIT200 and MEM18]
Isotype	IgG2a, IgG1
Buffer	PBS pH 7.2, 1% BSA, 0.05% NaN3

Form	FITC and PE
Formula	PBS pH 7.2, 1% BSA, 0.05% NaN3
Application Notes	<p>Staining Procedure Direct Immunofluorescence (Staining Procedure)</p> <p>Nordic-MUBio fluorochrome labeled antibodies are designed for use with either whole blood or isolated mononuclear cell (MNC) preparations. Proposed staining procedure for whole blood in short: - For each sample add 50 µl of EDTA anti-coagulated blood to a 3-5 ml tube - Add 20 µl of the appropriate Nordic-MUBio monoclonal antibody conjugate - Incubate the tube for 15 minutes at 4C or at room temperature in the dark - Add 100 µl NM-LYSE (Cat.No. GAS-003) to each tube and incubate for 10 minutes at room temperature - Add 3-4 ml of distilled water and vortex, incubate for 5-10 minutes at room temperature - Centrifuge tube for 5 minutes at 300 g - Aspirate supernatant and resuspend pellet in 0.3 ml of sheath fluid - Analyze immediately or store samples at 2-8 C in the dark and analyze within 24 hours For "No-Wash protocol please refer to <a href="http://www.nordicmubio.com">www.nordicmubio.com</a> Proposed staining procedure for MNC in short: - Carefully add 20 µl antibody conjugate and 50-100 µl MNC to the bottom of a tube - Vortex at low speed for 1-2 seconds - Incubate for 15-30 minutes at 2-8C or at room temperature - Centrifuge tubes for 5 minutes at 300 g - Remove supernatant, resuspend cells in 2-5 ml of phosphate buffered saline (PBS) and centrifuge cells again for 5 minutes at 300 g - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1 % formaldehyde and store them at 2-8C in the dark. Analyze fixed cells within 24 hours</p>