

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Bovine IgG (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GAB/IGG(H+L)/FITC

| | |
|----------------------------|---|
| Article Name | Goat anti Bovine IgG (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAB/IGG(H+L)/FITC |
| Supplier Catalog Number | GAB/IgG(H+L)/FITC |
| Alternative Catalog Number | NMB-GAB/IGG(H+L)/FITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Bovine |
| Conjugation | FITC |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | Human, Porcine |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes some reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in bovine serum, as tested by imm... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|---|
| Buffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. It is reconstituted by adding 2 ml sterile distilled water, spun down to remove insoluble particles, divided into sma |
| Source | Purified normal IgG isolated from pooled bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Form | FITC |
| Formula | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |