

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Chicken IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GACH/IGM(FC)

Article Name	Goat anti Chicken IgM (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GACH/IGM(FC)
Supplier Catalog Number	GACH/IgM(Fc)
Alternative Catalog Number	NMB-GACH/IGM(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Gallus
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal chicken serum a single precipitin line is obtained which show...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an
Source	Highly purified normal IgM isolated from pooled chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l in a 3 mm diameter center well and 2 $\mu$ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.