

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Duck IgG IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GADU/IG

| | |
|----------------------------|---|
| Article Name | Goat anti Duck IgG IgA IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GADU/IG |
| Supplier Catalog Number | GADu/Ig |
| Alternative Catalog Number | NMB-GADU/IG |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Aves |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgG+IgM+IgA (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins. It does not react with any non-Ig protein in duck serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techn... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|---|
| Buffer | Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat seru |
| Source | Purified normal Ig fractions isolated from pooled duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells. |